Neuroscience Letters, 169 (1994) 228–230
Elsevier Science Ireland Ltd.

AUTHOR INDEX

Agnati, L.F., see Fuxe, K., 25
Angelin, B., see Lannfelt, L., 175
Armstrong, R.A., Differences in β-amyloid (β/A4) deposition in human patients with Down's syndrome and sporadic Alzheimer's disease, 133
Artero, C., Fasolo, A. and Franzoni, M.F., Multiple sources of the pituitary pars intermedia innervation in amphibians: a DiI retrograde tract-tracing study, 163
Azzaro, A.J., see Gabr, R.W., 63

Backhovens, H., see Van Broeckhoven, C., 179
Baldissera, F., Loreto, S.D., Florio, T. and Scarnati, E., Short-latency excitation of hindlimb motoneurons induced by electrical stimulation of the pontomesencephalic tegmentum in the rat, 13
Barber, C.M., see Harrison, J.K., 85
Barillot, J.-C., see Frugière, A., 191
Bastianelli, E. and Pochet, R., Distribution of calmodulin, calbindin-D28k and calretinin among rat olfactory nerve bundles, 223
Basun, H., see Lannfelt, L., 175
Bates, F., see Donoso, M.V., 59
Beckmann, H., see Künig, G., 219
Berglund, L., see Lannfelt, L., 175
Biggs, C.S., see Whitton, P.S., 215
Birkle, D.L., see Gabr, R.W., 63
Bliss, T.V.P., see Fazeli, M.S., 77
Boissl, K.W., see Janetzky, B., 126
Brammer, M.J., see Romano-Silva, M.A., 90
Breen, K., see Fazeli, M.S., 77
Bulat, M., see Strikić, N., 159
Burton, M.D., Hitzig, B.M. and Johnson, D.C., Viability of the neonatal rat isolated brainstem preparation by ^{31}P MRS, 31

Cao, X., see Phillis, J.W., 188
Caravaglios, G., see Ferraro, G., 93
Carlton, S.M., see Steel, J.H., 81
Chadi, G., see Fuxe, K., 25
Chen, A.C., see Waite, J.J., 154
Cherubini, E., see Hosokawa, Y., 167
Chung, J.M., see Steel, J.H., 81
Cowburn, P.J. and Payne, A.P., Androgens and indoleamines interact to control sexual dimorphisms in the rat spinal cord, 101
Crockett, D.P., see Foschini, D.R., 21
Crook, R., see Van Broeckhoven, C., 179
Crossman, A.R., see Griffiths, P.D., 68
Cruts, M., see Van Broeckhoven, C., 179
Czyrak, A., see Pilc, A., 171

Decchi, B., see Rossi, A., 141
Deckert, J., see Künig, G., 219
Donoso, M.V., Bates, F., Montiel, J. and Huidobro-Toro, J.P., Adenosine 5″-triphosphate (ATP), the neurotransmitter in the prostatic portion of the longitudinal muscle layer of the rat vas deferens, 59
Dubourg, P., see Magoul, R., 149

Egger, M.D., see Foschini, D.R., 21
Eggertsen, G., see Lannfelt, L., 175
Errington, M.L., see Fazeli, M.S., 77

Fasolo, A., see Artero, C., 163
Fazeli, M.S., Breen, K., Errington, M.L. and Bliss, T.V.P., Increase in extracellular NCAM and amyloid precursor protein following induction of long-term potentiation in the dentate gyrus of anaesthetized rats, 77
Felix, D., Wu, G.-Y. and Wang, S.-R., GABA as an inhibitory transmitter in the pigeon isthmo-tectal pathway, 212
Ferraro, G., Sardo, P., Sabatino, M., Caravaglios, G. and La Grutta, V., Anticonvulsant activity of the noradrenergic locus coeruleus system: role of beta mediation, 93
Florio, T., see Baldissera, F., 13
Foschini, D.R., Kestler, A.M., Egger, M.D. and Crockett, D.P., The up-regulation of trkA and trkB in dorsal column astrocytes following dorsal rhizotomy, 21
Fowler, L.J., see Whitton, P.S., 215
Frankiewicz, T., see Pilc, A., 171
Franzoni, M.F., see Artero, C., 163
Fratiglioni, L., see Lannfelt, L., 175
Friedrich, R.W., Quigley, P.A., Srivastava, M., Skerrett, M. and Mercier, A.J., Temperature dependence of synaptic modulation by a FMRFamide-related neuropeptide in crayfish, 56
Frugière, A. and Barillot, J.-C., The medullary rostral ventrolateral pressor region: an electrophysiological study in decerebrate rat, 191
Fukuuchi, Y., see Isozumi, K., 203
Funada, M., Suzuki, T. and Misawa, M., The role of dopamine D_1-receptors in morphine-induced hyperlocomotion in mice, 1
Fuxe, K., Tinner, B., Chadi, G., Härfstrand, A. and Agnati, L.F., Evidence for a regional distribution of hyaluronic acid in the rat brain using a highly specific hyaluronic acid recognizing protein, 25

Gabr, R.W., Gladfelter, W.E., Birkle, D.L. and Azzaro, A.J., In vivo microdialysis of corticotropin releasing factor (CRF): calcium dependence of depolarization-induced neurosecretion of CRF, 63
Geiger, J.D., see Gu, J.G., 185
Gladfelter, W.E., see Gabr, R.W., 63
Goel, R., Mishra, O.P. and Papadopoulos, M.D., Effect of dithiothreitol on lipid peroxidation induced modification of NMDA receptor in fetal guinea pig brain, 109
Gomez, M.V., see Romano-Silva, M.A., 90
Greenamyre, J.T., see Porter, R.H.P., 105
Griffiths, P.D., Perry, R.H. and Crossman, A.R., A detailed anatomical analysis of neurotransmitter receptors in the putamen and caudate in Parkinson's disease and Alzheimer's disease, 68
Gu, J.G. and Geiger, J.D., Effects of diadenosine polyphosphates on sodium nitroprusside-induced soluble guanylate cyclase activity in rat cerebellum, 185

Hahn, S.J., Yoon, S.H., Sim, S.S., Rhie, D.J., Jo, Y.H. and Kim, M.S., Stimulatory effects of the central amygdaloid nucleus on pancreatic exocrine secretion in rats, 43
Hao, J.-X. and Wiesenfeld-Hallin, Z., Variability in the occurrence of ongoing discharges in primary afferents originating in the neuroma after peripheral nerve section in different strains of rats, 119
Hardy, J., see Van Broeckhoven, C., 179
Hardy, S.G.P., Anatomical data supporting the concept of prefrontal influences upon hypothalamo-medullary relays in the rat, 17
Härfstrand, A., see Fuxe, K., 25

Harrison, J.K., Barber, C.M. and Lynch, K.R., cDNA cloning of a G-protein-coupled receptor expressed in rat spinal cord and brain related to chemokine receptors, 85
Hartmann, J., see Künig, G., 219
Hatanaka, H., see Kojima, M., 47
Hauck, S., see Janetzky, B., 126
Heinsen, H., see Künig, G., 219
Henneguelle, E., see Pratt, J., 114
Hitzig, B.M., see Burton, M.D., 31
Homma, I., see Kanamaru, M., 35
Honmou, O., see Kocsis, J.D., 181
Hosokawa, Y., Sandri, G., Panfili, E. and Cherubini, E., Characterization of a voltage-dependent anionic channel in fused synaptosomes isolated from rat hippocampi, 167
Houlden, H., see Van Broeckhoven, C., 179
Huidobro-Toro, J.P., see Donoso, M.V., 59

Ikeuchi, T., see Kojima, M., 47
Inouye, T., Shinosaki, K., Iyama, A., Matsumoto, Y., Toi, S. and Ishihara, T., Potential flow of frontal midline theta activity during a mental task in the human electroencephalogram, 145
Ishihara, T., see Inouye, T., 145
Isozumi, K., Fukuuchi, Y., Takeda, H. and Itoh, Y., Cerebral ischemia-induced amplification phenomenon of somatosensory evoked potentials in cats, 203
Itoh, Y., see Isozumi, K., 203
Iwase, M., see Kanamaru, M., 35
Iyama, A., see Inouye, T., 145

Janetzky, B., Hauck, S., Youdim, M.B.H., Riederer, P., Jellinger, K., Pantucek, F., Zöchling, R., Boissl, K.W. and Reichmann, H., Unaltered aconitase activity, but decreased complex I activity in substantia nigra pars compacta of patients with Parkinson's disease, 126
Jellinger, K., see Janetzky, B., 126
Jo, Y.H., see Hahn, S.J., 43
Johanson, C.E. and Preston, J.E., Potassium efflux from infant and adult rat choroid plexuses: effects of CSF anion substitution, *N*-ethylmaleimide and Cl transport inhibitors, 207
Johnson, D.C., see Burton, M.D., 31

Kah, O., see Magoul, R., 149
Kanamaru, M., Iwase, M. and Homma, I., Central mediation of tracheal pressure through H_1 and H_2 receptors after histamine administration to the fourth ventricle of anesthetized rabbit, 35
Kanner, B.I., see Rauen, T., 137
Kasai, H., see Mori, A., 199
Katoh, K., see Koshimoto, H., 73
Kawata, M., see Yuri, K., 5
Kestler, A.M., see Foschini, D.R., 21
Kim, M.S., see Hahn, S.J., 43
Klarica, M., see Strikić, N., 159
Kocsis, J.D. and Honmou, O., Gabapentin increases GABA-induced depolarization in rat neonatal optic nerve, 181
Kojima, M., Ikeuchi, T. and Hatanaka, H., Nerve growth factor induces *trk*A mRNA expression in cultured basal forebrain cholinergic neurons from 17-day fetal rats, 47
Koshimoto, H., Katoh, K., Yoshihara, Y., Nemoto, Y. and Mori, K., Immunohistochemical demonstration of embryonic expression of an odor receptor protein and its zonal distribution in the rat olfactory epithelium, 73
Künig, G., Hartmann, J., Niedermeyer, B., Deckert, J., Ransmayr, G., Heinsen, H., Beckmann, H. and Riederer, P., Excitotoxins L-β-oxalyl-amino-alanine (L-BOAA) and 3,4,6-trihydroxyphenylalanine (6-OH-DOPA) inhibit [^{3}H]α-amino-3-hydroxy-5-methyl-4-isoxazole-propionic acid (AMPA) binding in human hippocampus, 219

La Grutta, V., see Ferraro, G., 93
Laduron, P.M., see Pratt, J., 114
Lannfelt, L., Lilius, L., Nastase, M., Viitanen, M., Fratiglioni, L., Eggertsen, G., Berglund, L., Angelin, B., Linder, J., Winblad, B. and Basun, H., Lack of association between apolipoprotein E allele ε4 and sporadic Alzheimer's disease, 175
Lappi, D.A., see Waite, J.J., 154
Legutko, B., see Pilc, A., 171
León-Olea, M., see Sánchez-Alvarez, M., 51
Lilius, L., see Lannfelt, L., 175
Linder, J., see Lannfelt, L., 175
Loreto, S.D., see Baldissera, F., 13
Lucas-Heron, B., Schmitt, N. and Ollivier, B., Mdx mouse skeletal muscle: could a mitochondrial factor be responsible for the absence of progressive necrosis?, 97
Luo, L. and Wiesenfeld-Hallin, Z., Genetic factors may influence the development of spinal reflex hyperexcitability following sciatic nerve section in the rat, 122
Lynch, K.R., see Harrison, J.K., 85

Magoul, R., Dubourg, P., Kah, O. and Tramu, G., Tachykinergic synaptic inputs to neurons of the medial preoptic region which project to the rat arcuate nucleus, 149
Marcotte, E.R., Sullivan, R.M. and Mishra, R.K., Striatal G-proteins: effects of unilateral 6-hydroxydopamine lesions, 195
Martin, J.J., see Van Broeckhoven, C., 179
Martínez-Lorenzana, G., see Sánchez-Alvarez, M., 51
Matsumoto, Y., see Inouye, T., 145
Mercier, A.J., see Friedrich, R.W., 56
Miller, J.W., see Torack, R.M., 9
Misawa, M., see Funada, M., 1
Mishra, O.P., see Goel, R., 109
Mishra, R.K., see Marcotte, E.R., 195
Miyashita, Y., see Mori, A., 199
Montiel, J., see Donoso, M.V., 59
Moraes-Santos, T., see Romano-Silva, M.A., 90
Mori, A., Takahashi, T., Miyashita, Y. and Kasai, H., Quantal properties of S-type glutamatergic synaptic input to the striatal medium spiny neuron from neonate rat, 199
Mori, K., see Koshimoto, H., 73
Mori, N. and Watanabe, M., Comparison of effects of bilateral injections of bicuculline and muscimol into the caudate-putamen of amygdaloid-kindled rats, 129

Na, H.S., see Steel, J.H., 81
Nastase, M., see Lannfelt, L., 175
Nemoto, Y., see Koshimoto, H., 73
Niedermeyer, B., see Künig, G., 219

Ollivier, B., see Lucas-Heron, B., 97

Panfili, E., see Hosokawa, Y., 167
Pantucek, F., see Janetzky, B., 126
Papadopoulos, M.D., see Goel, R., 109
Payne, A.P., see Cowburn, P.J., 101
Pellicer, F., see Sánchez-Alvarez, M., 51
Perry, R.H., see Griffiths, P.D., 68

Phillis, J.W., Sen, S. and Cao, X., Amflutizole, a xanthine oxidase inhibitor, inhibits free radical generation in the ischemic/reperfused rat cerebral cortex, 188
Pilc, A., Legutko, B., Frankiewicz, T. and Czyrak, A., Trans-(±)-1-amino-1,3-cyclopentanedicarboxylate (trans-ACPD) stimulates cAMP accumulation in rat cerebral cortical slices but not in glial or neuronal cell cultures, 171
Pochet, R., see Bastianelli, E., 223
Polak, J.M., see Steel, J.H., 81
Porter, R.H.P. and Greenamyre, J.T., Inhibition of autoradiographic MK-801 binding by an endogenous factor, 105
Pratt, J., Roux, M., Henneguelle, E., Stutzmann, J.-M. and Laduron, P.M., Neuroprotective effects of colchicine in the gerbil model of cerebral ischaemia, 114
Preston, J.E., see Johanson, C.E., 207

Quasthoff, S., A mechanosensitive K^+ channel with fast-gating kinetics on human axons blocked by gadolinium ions, 39
Quigley, P.A., see Friedrich, R.W., 56

Ransmayr, G., see Künig, G., 219
Rauen, T. and Kanner, B.I., Localization of the glutamate transporter GLT-1 in rat and macaque monkey retinae, 137
Reichmann, H., see Janetzky, B., 126
Rhie, D.J., see Hahn, S.J., 43
Ribeiro-Santos, R., see Romano-Silva, M.A., 90
Richards, D.A., see Whitton, P.S., 215
Riederer, P., see Künig, G., 219
Riederer, P., see Janetzky, B., 126
Romano-Silva, M.A., Ribeiro-Santos, R., Gomez, M.V., Moraes-Santos, T. and Brammer, M.J., Tityustoxin-mediated Na^+ influx is more efficient than KCl depolarisation in promoting Ca^{2+}-dependent glutamate release from synaptosomes, 90
Rossi, A., Zalaffi, A. and Decchi, B., Heteronymous recurrent inhibition from gastrocnemius muscle to soleus motoneurones in humans, 141
Roux, M., see Pratt, J., 114

Sabatino, M., see Ferraro, G., 93
Sánchez-Alvarez, M., León-Olea, M., Talavera, E., Pellicer, F., Sánchez-Islas, E. and Martínez-Lorenzana, G., Distribution of NADPH-diaphorase in the perioesophageal ganglia of the snail, *Helix aspersa*, 51
Sánchez-Islas, E., see Sánchez-Alvarez, M., 51
Sandri, G., see Hosokawa, Y., 167
Sardo, P., see Ferraro, G., 93
Scarnati, E., see Baldissera, F., 13
Schmitt, N., see Lucas-Heron, B., 97
Sen, S., see Phillis, J.W., 188
Shinosaki, K., see Inouye, T., 145
Sim, S.S., see Hahn, S.J., 43
Skerrett, M., see Friedrich, R.W., 56
Srivastava, M., see Friedrich, R.W., 56
Steel, J.H., Terenghi, G., Chung, J.M., Na, H.S., Carlton, S.M. and Polak, J.M., Increased nitric oxide synthase immunoreactivity in rat dorsal root ganglia in a neuropathic pain model, 81
Strikić, N., Klarica, M., Vladić, A. and Bulat, M., Effect of active transport on distribution and concentration gradients of [^{3}H]benzylpenicillin in the cerebrospinal fluid, 159
Stutzmann, J.-M., see Pratt, J., 114
Sullivan, R.M., see Marcotte, E.R., 195
Suzuki, T., see Funada, M., 1

Takahashi, T., see Mori, A., 199
Takeda, H., see Isozumi, K., 203
Talavera, E., see Sánchez-Alvarez, M., 51
Terenghi, G., see Steel, J.H., 81
Thal, L.J., see Waite, J.J., 154
Tinner, B., see Fuxe, K., 25
Toi, S., see Inouye, T., 145
Torack, R.M. and Miller, J.W., Immunoreactive changes resulting from dopaminergic denervation of the dentate gyrus of the rat hippocampal formation, 9
Tramu, G., see Magoul, R., 149

Van Broeckhoven, C., Backhovens, H., Cruts, M., Martin, J.J., Crook, R., Houlden, H. and Hardy, J., APOE genotype does not modulate age of onset in families with chromosome 14 encoded Alzheimer's disease, 179
Viitanen, M., see Lannfelt, L., 175
Vladić, A., see Strikić, N., 159

Waite, J.J., Wardlow, M.L., Chen, A.C., Lappi, D.A., Wiley, R.G. and Thal, L.J., Time course of cholinergic and monoaminergic changes in rat brain after immunolesioning with 192 IgG-saporin, 154
Wang, S.-R., see Felix, D., 212
Wardlow, M.L., see Waite, J.J., 154
Watanabe, M., see Mori, N., 129
Whitton, P.S., Richards, D.A., Biggs, C.S. and Fowler, L.J., *N*-Methyl-D-aspartate receptors modulate extracellular 5-hydroxy- tryptamine concentration in rat hippocampus and striatum in vivo, 215
Wiesenfeld-Hallin, Z., see Luo, L., 122
Wiesenfeld-Hallin, Z., see Hao, J.-X., 119
Wiley, R.G., see Waite, J.J., 154
Winblad, B., see Lannfelt, L., 175
Wu, G.-Y., see Felix, D., 212

Yoon, S.H., see Hahn, S.J., 43
Yoshihara, Y., see Koshimoto, H., 73
Youdim, M.B.H., see Janetzky, B., 126
Yuri, K. and Kawata, M., Estrogen affects calcitonin gene-related peptide- and methionine-enkephalin-immunoreactive neuron in the female rat preoptic area, 5

Zalaffi, A., see Rossi, A., 141
Zöchling, R., see Janetzky, B., 126